## Diazepam and Nordiazepam

Order Name:DIAZEPAMTest Number:3611375Revision Date:10/18/2017

TEST NAME			METHODOLOGY	LOINC CODE	
Diazepam and Nordiazepam			Liquid Chromatography/Tandem Mass Spectrometry		
SPECIMEN REQUIRE	MENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment	
Preferred	1 mL (0.4 mL)	Serum	Clot Activator (Red Top, No-Gel)	Refrigerated	
Instructions	<ul> <li>Specimen Type: Red-top tube, lavender-top (EDTA) tube OR green-top (heparin) tube. DO NOT USE A GEL-BARRIER TUBE. The use of gelbarrier tubes is not recommended due to slow absorption of the drug by the gel. Depending on the specimen volume and storage time, the decrease in drug level due to absorption may be clinically significant.</li> <li>Specimen Storage: Refrigerate</li> <li>Specimen Collection: Transfer separated serum or plasma to a plastic transport tube. Oral: peak; one hour after dose; I.V.: peak: 15 minutes after dose.</li> <li>Specimen Stability: Ambient: 1 day, Refrigerated : 14 days, Frozen: 14 days</li> </ul>		olume and storage time, the decrease		

GENERAL INFORMATION		
Expected TAT	2-4 Days	
Performing Labcorp Test Code	007989	
Notes	Labcorp Test Code: 007989	
CPT Code(s)	80299	
Lab Section	Reference Lab	

Service provided by
Labcorp Oklahoma, Inc.
All Rights Reserved. © 2003 - 2024