

BCR/ABL Gene Rearrangement Quantitative by PCR

Order Name: **BCRABL PCR**
Test Number: 9101855
Revision Date: 10/01/2022

TEST NAME	METHODOLOGY	LOINC CODE
BCR/ABL Gene Rearrangement Quantitative by PCR	Polymerase Chain Reaction	

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	5 mL (3 mL)	Whole Blood	EDTA (Lavender Top)	Refrigerated
Instructions	<p>For Best results: Send specimens for testing ASAP. Specimens not going to be tested immediately should be stored refrigerated, specimens kept at room temperature will degrade faster than those kept refrigerated. Frozen samples will be rejected.</p> <p>CRITICAL: Do not open or share tube!</p> <p>Stability: Ambient 1 day(s), Refrigerated 2 day(s), Frozen: Unacceptable.</p>			

GENERAL INFORMATION	
Testing Schedule	Monday and Thursday
Expected TAT	Within 7 day(s)
Clinical Use	The bcr/abl rearrangement is detected in 90 to 95% of CML, some acute lymphocytic leukemia (ALL), and, rarely, in acute myelogenous leukemia (AML). Diagnose chronic myelogenous leukemia (CML) in the presence or absence of Philadelphia chromosome. Determine prognosis & relapse. Also used to identify acute lymphocytic leukemia (ALL) patients who have a Philadelphia chromosome.
Notes	The bcr/abl gene rearrangement is observed in CML, ALL, and AML. A negative result indicates fewer than 1 leukemic cell per 10,000 normal cells. This test detects only the bcr/abl translocation. It will not detect other translocations that may appear in the terminal phase of CML.
CPT Code(s)	81206, 81207, G0452-26 (can be a combination or all 3 codes)
Lab Section	Reference Lab