

HPV Only (ThinPrep Pap Collection)

Order Name: **HPV Only**

Test Number: 6910103

Revision Date: 10/10/2018

| TEST NAME | METHODOLOGY | LOINC CODE |
|---------------------------|-------------|------------|
| HPV Screen Interpretation | | |

| SPECIMEN REQUIREMENTS | | | | |
|-----------------------|---|---------------|------------------------------|-----------------------|
| Specimen | Specimen Volume (min) | Specimen Type | Specimen Container | Transport Environment |
| Preferred | Cervical brush | PAP specimen | ThinPrep PreservCyt solution | Room Temperature |
| Instructions | <p>Stability: Specimen must be transferred from ThinPrep PAP vial into an Aptima Specimen Transfer tube within 30 days. Once transferred, the specimen is stable in the Aptima vial for 30 days refrigerated.</p> <p>Patient information The patient should be tested 2 weeks after the first day of her last menstrual period, and definitely not when she is menstruating. The patient should not use vaginal medication, vaginal contraceptives, or douches for 48 hours before the exam. The patient should refrain from intercourse 48 hours prior to the exam.</p> <p>Preferred Specimens are samples from the female genital tract for Pap testing should be obtained with a cytobrush/spatula combination or a broom collection device and rinsed into a ThinPrep Pap® vial containing PreservCyt® solution.</p> <p>Specimen collection preparation Prepare the speculum. For patients without physical or physiological need for lubricant, use lukewarm water to warm and lubricate the speculum. When necessary, sparingly apply carbomer-free lubricant on the exterior of the speculum blades. Do not use an excessive amount of lubricant jelly to lubricate the speculum. Remove excess mucus or other discharge present before taking the sample. This should be gently removed with ring forceps holding a folded gauze pad. Remove inflammatory exudate from the cervical canal before taking the sample. Remove by placing a dry 2-by-2-inch piece of gauze over the cervix and peeling it away after it absorbs the exudate or by using a dry procto swab or Scopette swab. The cervix should not be cleaned by washing with saline or it may result in a relatively acellular specimen. The sample should be obtained before the application of acetic acid.</p> | | | |

| GENERAL INFORMATION | |
|---------------------|-----------------------|
| Testing Schedule | Mon-Sat |
| Expected TAT | 1-2 days after set-up |
| CPT Code(s) | 87624 |
| Lab Section | Molecular Diagnostic |