

Ct/GC/Tv NAA+M genitalium Urine

Order Name: **Ct/GC/Tv M.gen Ur**
Test Number: **5195318**
Revision Date: **05/01/2026**

TEST NAME	METHODOLOGY	LOINC CODE
Ct/GC/Tv NAA+M genitalium Urine	Nucleic acid amplification (NAA)	

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	30 mL (5 mL)	Urine (See Inst.)	APTIMA Urine Transport Tube	Room Temperature
Instructions	Specimen: First-void urine in Aptima® urine transport Volume: 5 mL or 30 mL urine collection Container: Aptima® urine transport Collection: Urine specimen: The patient should not have urinated for at least one hour prior to specimen collection. Direct patient to provide a first-catch urine (approximately 20 to 30 mL of the initial urine stream) into a urine collection cup free of any preservatives. Collection of larger volumes of urine may result in specimen dilution that may reduce test sensitivity; lesser volumes may not adequately rinse organisms into the specimen. Female patients should not cleanse the labial area prior to providing the specimen. Add urine to the Aptima® Combo 2 urine collection device. The final volume must be between the two black lines on the device (about 2 mL). Storage Instructions: Maintain specimen at room temperature or refrigerate (2°C to 30°C). Stability Room temperature 30 days (stability provided by manufacturer or literature reference)			

GENERAL INFORMATION			
Expected TAT	3-4 days		
Notes			
	RESULT CODE	RESULT NAME	LOINC CODE
	6907667	Mycoplasma genitalium Urine	88226-6
	6910159	Trichomonas Urine	70167-2
	6910151	Chlamydia Urine	42931-6
	6910155	Gonorrhea Urine	60256-5
Lab Section	Molecular Diagnostic		